

Veterinär-Anatomisches Institut
der Vetsuisse-Fakultät Universität Zürich
Direktor: Prof. Dr. med. vet. Alois Boos

Arbeit unter wissenschaftlicher Betreuung von Prof. Dr. Mariusz P. Kowalewski

***In vitro* decidualisation of canine uterine stromal cells**

Inaugural Dissertation

zur Erlangung der Doktorwürde der
Vetsuisse-Fakultät Universität Zürich

vorgelegt von

Ewa Kautz

Tierärztin
aus Gostyń, Polen

genehmigt auf Antrag von

Prof. Dr. Mariusz P. Kowalewski, Referent

Zürich 2015

Veterinär-Anatomisches Institut
der Vetsuisse-Fakultät Universität Zürich
Direktor: Prof. Dr. med. vet. Alois Boos

Arbeit unter wissenschaftlicher Betreuung von Prof. Dr. Mariusz P. Kowalewski

***In vitro* decidualisation of canine uterine stromal cells**

Inaugural Dissertation

zur Erlangung der Doktorwürde der
Vetsuisse-Fakultät Universität Zürich

vorgelegt von

Ewa Kautz

Tierärztin
aus Gostyń, Polen

genehmigt auf Antrag von

Prof. Dr. Mariusz P. Kowalewski, Referent

Zürich 2015

TABLE OF CONTENTS

1	SUMMARY	5
2	ZUSAMMENFASSUNG	6
3	AUTHORS AND CORRESPONDENCE	7
4	ABSTRACT	8
	Background	8
	Methods	8
	Results	8
	Conclusion	9
5	INTRODUCTION.....	10
6	MATERIAL AND METHODS.....	13
6.1	COLLECTION OF TISSUES AND ISOLATION OF CELLS	13
6.2	LIVE CELL IMAGING MICROSCOPY	14
6.3	IMMUNOCYTOCHEMISTRY AND IMMUNOHISTOCHEMISTRY	15
6.4	ELECTRON MICROSCOPY ANALYSIS	16
6.5	RNA ISOLATION, REVERSE TRANSCRIPTION (RT), SEMI-QUANTITATIVE (TAQMAN) PCR AND DATA EVALUATION.....	17
6.6	WESTERN BLOT ANALYSIS	19
7	RESULTS.....	21
7.1	CHARACTERISATION AND <i>IN VITRO</i> DECIDUALISATION OF ISOLATED CANINE UTERINE STROMAL CELLS.....	21
7.2	SUBCELLULAR MORPHOLOGICAL CHARACTERISATION OF CANINE DECIDUAL CELLS DURING <i>IN VITRO</i> DECIDUALISATION.	22
7.3	EXPRESSION OF SELECTED GENES IN CANINE UTERINE PRIMARY STROMAL CELLS DURING <i>IN VITRO</i> DECIDUALISATION.....	22
8	DISCUSSION	25
9	FIGURES AND LEGENDS	29
9.1	TABLE 1	29
9.2	FIGURE 1.....	30
9.3	FIGURE 2.....	31
9.4	FIGURE 3.....	32
9.5	FIGURE 4.....	33
9.6	FIGURE 5.....	34
9.7	FIGURE 6.....	35
9.8	FIGURE 7.....	36
9.9	FIGURE 8.....	37

9.10 SUPPLEMENT 1	38
10 FUNDING	39
11 REFERENCES.....	40
12 ACKNOWLEDGEMENTS.....	45
13 CURRICULUM VITAE	46

1 Summary

Uterine recognition of the dog embryo and establishing of canine pregnancy is still poorly understood. The intimate embryo-maternal cross-talk leads to strong structural and functional remodelling of the uterus called decidualisation, with maternal stromal cells transforming into decidual cells expressing variety of factors facilitating implantation. Understanding this process is essential for understanding canine reproductive physiology. Here, the capability of canine uterine stromal cells to decidualise *in vitro*, thereby serving as a model of decidualisation, was investigated using a dbcAMP- mediated approach. Tissue material from healthy bitches in early dioestrus was used. Cells were characterized by differential staining, nearly 100 % of which were vimentin-positive. Scanning and transmission electron microscope analyses were applied, and morphological changes were recorded with a live cell imaging microscope. Several decidualisation markers were investigated. Cultured stromal cells acquired characteristics of decidual cells when incubated with 0.5 mM dbcAMP for 72 h. Their shape changed from elongated to rounded with higher numbers of mitochondria and secretory follicles, and increased proliferation rate. Decidualisation markers, hormone receptors, members of prostaglandin system were elevated in decidualised cells. Herewith, a valuable *in vitro* model has been developed allowing the study of the underlying molecular and endocrine mechanisms of canine decidualisation.

2 Zusammenfassung

Die Embryo-maternale Trächtigkeitserkennung bei der Hündin bleibt weitgehend ungeklärt. Die strukturelle und funktionelle Umgestaltung des Uterusgewebes führt zur Umwandlung der uterinen Stromazellen in die sog. Dezidualisierung dieser Zellen. Die entstandenen Dezidualzellen exprimieren mehrere Faktoren, welche die Implantation ermöglichen. Diesen Vorgang zu verstehen scheint essentiell für das Verständnis der kaninen Reproduktionsphysiologie. Hierfür wurden die uterinen Stromazellen unter der Verwendung von dbcAMP *in vitro* dezidualisiert. Gewebe von gesunden Hündinnen im frühen Diöstrus wurde verwendet. Differentialfärbung wurde angewendet, wobei nahezu 100% Zellen vimentin-positiv waren. Raster- und transmissionsmikroskopische Analysen wurden angewandt und morphologische Veränderungen wurden mit einem Live Cell imaging Mikroskop dokumentiert. Mehrere Dezidualisationsmarkern wurden untersucht. Nach 72 Stunden Inkubation in Zellkultur mit 0.5mM dbcAMP erwarben die Stromazellen die Eigenschaften der Dezidualzellen. Ihre Gestalt wurde von länglich zu rundlich mit erhöhter Anzahl an Mitochondrien und sekretorischen Follikeln, zudem zeigten sie eine erhöhte Proliferationsrate. Dezidualisationmarker, Hormonrezeptoren und Mitglieder des Prostaglandinsystems waren erhöht in den Dezidualzellen. Hiermit wurde ein funktionelles *in vitro* Modell entwickelt, welches weitergehende Untersuchungen der molekularen- und endokrinologischen Mechanismen der caninen Dezidualisierung ermöglicht.

3 Authors and correspondence

***In vitro* decidualisation of canine uterine stromal cells.**

Authors: Ewa Kautz¹, Paula de Carvalho Papa², Iris M. Reichler³, Aykut Gram¹, Alois Boos¹,
Mariusz P. Kowalewski¹

Published in:

Reproductive Biology and Endocrinology, 2015 Aug 5;13:85.

doi: 10.1186/s12958-015-0066-4.

E Kautz¹ : ewa.kautz@yahoo.com

PC Papa² : paula.papa1@gmail.com

IM Reichler³ : ireichler@vetclinics.uzh.ch

A Gram¹ : aykutgram@gmail.com

A Boos¹ : boosa@vetanat.uzh.ch

MP Kowalewski¹ : kowalewski1@yahoo.de

¹⁾ Institute of Veterinary Anatomy, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland.

²⁾ Sector of Anatomy, Department of Surgery, Faculty of Veterinary Medicine and Animal Sciences, University of Sao Paulo, Sao Paulo, Brazil.

³⁾ Section of Small Animal Reproduction, Clinic of Reproductive Medicine, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland.

Correspondence:

Prof. Dr. Mariusz P. Kowalewski

Institute of Veterinary Anatomy

Vetsuisse Faculty

University of Zurich

Winterthurerstrasse 260

CH-8057 Zurich, Switzerland

Tel.: 0041-44-6358784; Fax: 0041-44-6358943

Email: kowalewski1@yahoo.de or kowalewski@vetanat.uzh.ch

4 Abstract

Background

The uterine response to the presence of embryos is poorly understood in the domestic dog (*Canis familiaris*). The intimate embryo-maternal cross-talk, which begins following the hatching of blastocysts and embryo attachment leads to strong structural and functional remodelling of the uterus. A part of this process is decidualisation, comprising morphological and biochemical changes that result in formation of maternal stroma-derived decidual cells. These are an integral part of the canine *placenta materna*, which together with the maternal vascular endothelium are the only cells of the canine endotheliochorial placenta able to resist trophoblast invasion. These cells are also the only ones within the canine placenta expressing the progesterone receptor (PGR). Understanding the decidualisation process thus appears essential for understanding canine reproductive physiology.

Methods

Here, we investigated the capability of canine uterine stromal cells to decidualise *in vitro*, thereby serving as a canine model of decidualisation. A dbcAMP-mediated approach was chosen during a time course of 24 - 72 h. Tissue material from six ($n = 6$) healthy, dioestric bitches was used (approximately 2 weeks after ovulation). Cells were characterized by differential staining, nearly 100 % of which were vimentin-positive. Scanning and transmission electron microscope analyses were applied, and morphological changes were recorded with a live cell imaging microscope. Expression of several decidualisation markers was investigated.

Results

The *in vitro* cultured stromal cells acquired characteristics of decidual cells when incubated with 0.5 mM dbcAMP for 72 h. Their shape changed from elongated to rounded, while ultrastructural analysis revealed higher numbers of mitochondria and secretory follicles, and an increased proliferation rate. Elevated expression levels of IGF1, IGF2, PRLR and ER α

were observed in decidualised cells; PRL and ER β remained mostly below the detection limit, while PGR remained unaffected. The expression of smooth muscle α actin (α SMA), another decidualisation marker, was strongly induced. Among prostaglandin system members, levels of COX2 (PTGS2) and of PGE2-synthase (PTGES) were upregulated. Expression of the PGE2 receptors, PTGER2 and PTGER4, was clearly detectable.

Conclusion

An *in vitro* decidualisation model with canine uterine stromal cells was successfully established, allowing future, more detailed studies to be undertaken on the underlying molecular and endocrine mechanisms of canine decidualisation.

Keywords: *Domestic dog; Decidualisation; in vitro model*

5 Introduction

Successful establishment of pregnancy requires highly orchestrated interactions between embryonic and maternal uterine tissues, which undergo specific morphological and biochemical changes to establish the uterine milieu required for proper embryo development. In animal species exhibiting invasive types of placentation, *e.g.*, higher order primates and rodents which develop hemochorial placentas, or dogs and cats which exhibit endotheliochorial placentas, these restructuring changes are very intense and result in decidua formation. Reprogramming of the endometrial stromal compartment during decidualisation involves transformation of stromal cells into decidual cells, which are involved in coordinating the processes of embryo implantation, placenta formation and development of the conceptus [1, 2]. This specific tissue differentiation remains under the control of progesterone (P4) and oestrogens [3, 4]. Problems during decidualisation are frequent causes of implantation failures in humans [5], and may contribute to abnormal placentation, sometimes resulting in severe conditions such as formation of *Placenta accreta* which is characterised by excessive trophoblast invasion into the decidua [6]. In other species, such as dogs, the latter anomaly, *i.e.* exaggerated trophoblast invasion, can result in a condition known as subinvolution of placental sites (SIPS) [7].

In the domestic dog, the establishment and maintenance of pregnancy depends entirely on P4 secreted from Corpora lutea (CL) because there is no placental steroidogenesis in this species [8-10]. Furthermore, in the absence of pregnancy, the dog also lacks an endogenous luteolysin, which results in a similar luteal life span and circulating hormone profiles during pregnancy and in pseudopregnant bitches [11]. Displaying these unique features, reproductive function in the dog differs distinctly from other domestic animal species. Consequently, being devoid of an embryo-derived anti-luteolytic principle found in livestock, *e.g.*, cattle or pigs [12-14], the dog appears to be an interesting model for investigating alternative mechanisms

involved in the establishment of pregnancy in mammals. However, there is only little available information concerning the uterine milieu prior to and during implantation and placentation in dogs. Just recently [15], the effects exerted by free-floating embryos (days 10-12 after fertilisation) on the canine pre-implantation uterus were investigated, revealing increased expression of several decidualisation markers, *e.g.*, insulin-like growth factor (IGF) 2, prolactin receptor (PRLR) and oestrogen receptor α (ER α , ESR1). Besides being clearly detectable in the uterine epithelial compartments, their expression was also detected in stromal cells. Interestingly, in contrast to rodents or primates [16, 17], the capacity of the canine uterus to produce PRL was found to be very low [15]. Among the prostaglandin (PG) family members, especially the expression of PGE2-synthase (PTGES) and one of the respective PGE2 receptors, PTGER2, was significantly upregulated in response to the presence of free-floating embryos [15].

The embryo-induced uterine functional differentiation in the dog becomes very intense following implantation, *i.e.*, by days 17-18 after fertilisation and placentation [18, 19]. The latter event is characterised by strong, trophoblast-induced structural and functional remodelling of the endometrium, eventually leading to the formation of highly specialised canine decidual cells. Together with maternal vascular endothelial cells, canine decidual cells are able to resist the strong proteolytic activity of embryonic cells. Thus, their role in modulating trophoblast invasiveness, and thereby protecting maternal tissues from excessive degradation, can be assumed. Most importantly, these are also the only cells of the dog placenta expressing PGR and ER α (ESR1) and, thereby, must play a pivotal role in the maintenance of pregnancy [20, 21]. Interfering with their function, rather than their expression, *e.g.*, by applying a selective PGR blocker, will unequivocally lead to pre-term parturition/abortion by activating utero-placental PG synthesis [20]. Acting at the level of decidual cell-targeted expression of its receptor (OTR, OXTR), oxytocin appears to be an

important mediator in this process [22]. Nevertheless, the underlying endocrine mechanisms associated with the decidualisation process in the dog remain unknown.

In vitro decidualisation can be induced by several stimuli. One of these, cyclic AMP (cAMP), is not only an important mediator of P4-induced decidualisation [23], but acts as a stronger stimulus than P4 in inducing expression of decidualisation markers in endometrial stromal cells [24-26]. Here, aiming to understand decidualisation in the dog, the capability of canine uterine stromal cells to undergo this process *in vitro* was investigated. Cells were isolated from uteri of naturally oestrogenized, dioestrous dogs. Due to our limited access to the experimental material, and bearing in mind the above-mentioned strong decidualisation potential of cAMP, a cAMP-based protocol was applied. Then, specific morphological, ultrastructural and functional phenomena associated with the *in vitro* decidualisation process were identified and evaluated.

6 Material and Methods

6.1 Collection of tissues and isolation of cells

Uterine tissues were used from six (n=6) non-pregnant, clinically healthy bitches, submitted for routine ovariohysterectomy at the Section of Small Animal Reproduction, Clinic of Reproductive Medicine, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland. Surgeries were performed during early non-pregnant dioestrus, approximately 12-14 days after clinical signs of heat had ceased. Immediately after collection, uteri were washed with sterile phosphate-buffered saline (PBS) solution and separated from surrounding connective tissues. Uteri were slit longitudinally and the endometrial layers were mechanically separated. After washing with PBS, tissue fragments were enzymatically dissociated by stirring for 2h at 37°C in PBS containing 0.15% collagenase (Sigma-Aldrich Chemie GmbH, Buchs, CH). Subsequent removal of undissociated tissue fragments was achieved by filtering through a 75µm nylon strainer (BD Biosciences, Basel, CH). Following this, cells were washed three times in PBS with centrifugation steps in between (400xg, 10 min, 4°C). Afterwards, pellets were resuspended with culture medium (DMEM High Glucose, pH 7.2-7.4, with 10% heat inactivated FBS, containing 100U/ml penicillin and 100µg/ml streptomycin, and 1% ITS (Insulin-Transferrin-Selenium); all from Chemie Brunschwig AG, Basel, CH) and seeded into 150 ml culture flasks (Corning, Amsterdam, NL). Following 1h of incubation in a humidified incubator at 37°C under 5% CO₂ in air, needed for adhesion of fibroblast stromal cells and their separation from epithelial cells (differential adhesion time), the medium was replaced with fresh cell-free medium. The attached stromal cells were further cultured under the same conditions until confluence, which was reached after approximately 4-5 days. Cells after the first passage were used for further experiments. Then, after trypsinisation and harvesting, cells were seeded into 6-well plates at different concentrations: 4×10^4 per well for 24h and 2

$\times 10^4$ per well for 48h and 72h incubation, respectively. When seeded at less than 2×10^4 per well, cells easily attached but exhibited retarded growth, most likely due to the low density and lack of sufficient cell-to-cell contact. On the other hand, when cultured longer than 72h, stimulated cells became over-confluent. For immunocytochemistry, sterile glass cover slips were placed into the wells to allow the seeded cells to adhere to them. Following seeding, cells were further cultured under the above conditions for approximately 24h and then used for experiments. Prior to stimulation, cells were washed with PBS, thereby removing the serum-containing medium. Subsequently, cells were treated with N⁶,2'-O-Dibutyryladenosine-3',5'-cyclic monophosphate (dbcAMP) in stimulation medium (DMEM High Glucose, pH = 7.2-7.4; penicillin/streptomycin as above), containing 0.1% bovine serum albumin (BSA), in the following concentrations: 0mM dbcAMP for the non-treated control, or 0.1mM, 0.3mM or 0.5mM dbcAMP. In pilot experiments, serum-free medium was used for stimulation, which resulted in detachment of cells within 2h after stimulation. Inclusion of BSA at the indicated concentration of 0.1% proved to be the lowest threshold determined empirically in experiments comprising serial dilution of BSA, needed for cells to prevent their detachment (data not shown).

6.2 Live cell imaging microscopy

Cells seeded in 6-well plates were placed on a LeicaDMI 6000B fluorescence microscope equipped with a Leica DFC360FX camera and an automated stage for live cell imaging (allowing for simultaneous observation of cells undergoing *in vitro* decidualisation and comparison with non-treated controls), enclosed within a thermostatically-controlled cabinet. Stimulations for live cell imaging microscopy were done in duplicate as follows: non-treated controls and 0.5 mM dbcAMP. The focal planes were set for image capture, and images were taken with a 10x objective every 5min for 72h, resulting in a total of 864 cycles. Culture

conditions were: 20% O₂, 5% CO₂, at 37°C. The resulting time-lapse videos were displayed at a fast speed to provide an accelerated view of *in vitro* decidualisation of canine stromal cells. Representative videos of control and *in vitro* decidualised canine stromal cells are presented in the supplemental material. The density of cells was quantified at cycles 144 (12h), 288 (24h), 576 (48h) and 864 (72h) and was evaluated relative to the surface of the applied fields of view using the MCID Analysis Software (InterFocus Limited, Linton, UK). *N*-fold changes in relative target area were determined by applying one-way analysis of variance (ANOVA) followed by Dunnett's Multiple Comparison test. Student's *t*-test was applied to test the effect of treatment on cell density at the selected time points. Statistical tests were performed using the GraphPad 3.06 program (GraphPad Statistical Software, San Diego, CA, USA). Results are presented as *n*-fold changes in the relative target area, reflecting the density of cells as % of the surface.

6.3 Immunocytochemistry and immunohistochemistry

Immunofluorescence was used for staining canine uterine primary cells seeded in 6-well plates on sterile cover slips. Cells were fixed by adding formaldehyde to cell cultures at a final concentration of 2% for 10 min at 37°C. After fixation, each well was washed twice with ice-cold PBS and submitted to the staining procedure described previously [27]. Briefly: cover slips were washed twice with ice-cold PBST (PBS/0.25% Triton X) followed by antigen retrieval in 50mM glycine (Sigma-Aldrich) in PBS for 5 min, and two washing steps in PBST (5 min each). A 1% solution of goat serum diluted in PBST was used for blocking non-specific binding sites (30 min). Next, primary antibodies were applied for 2h. These were: monoclonal mouse anti-vimentin (dilution 1:100), affinity-purified polyclonal rabbit anti-cytokeratin (dilution 1:300) and monoclonal mouse anti-human smooth muscle alpha actin (α SMA) (dilution 1:100), all from from Dako North America, Inc., CA, USA. Negative

controls included: samples omitting primary antibody, and those where both primary and secondary antibodies were omitted (for the autofluorescence control). Fluoresceinated goat anti-rabbit IgG antibody FI-1000 (Vector Laboratories, Burlingame, CA, USA) and fluoresceinated goat anti-mouse IgG antibody FI-2000, were used as secondary antibodies and applied in PBST for 1h. Nuclear staining was achieved by adding 4',6-diamidino-2-phenylindole (DAPI) (Sigma-Aldrich) to the secondary antibody solution. Cells were post-fixed with 2% formaldehyde. Glycergel® (Dako North America) was used for mounting cover slips on microscope slides.

Uterine cross-sections 2-3µm thick from early pregnant bitches, pre-implantation stages of pregnancy (days 10-12 after fertilisation; n=3) and utero-placental compartments from mid-pregnant dogs (35-40 days of pregnancy; n=3), were immunohistochemically stained for αSMA with our standard immunoperoxidase method [28, 29]. The primary antibody was the same as for the immunocytochemistry and used at 1:100 dilution. The secondary antibody was horse anti-mouse IgG BA-2000 (dilution 1:100; Vector Laboratories). The Vectastain ABC kit (Vector Laboratories) was used for enhancing signals. Peroxidase detection was done with the Liquid DAB+ substrate kit (Dako Schweiz AG, Baar, CH). Negative controls included: omitting the primary antibody, and an isotype control, rabbit IgG irrelevant antibodies I-1000 from Vector Laboratories used at the same concentration as the primary antibody (not shown).

6.4 Electron microscopy analysis

Previously described protocols for conventional transmission electron microscopy (TEM) and scanning electron microscopy (SEM) were used [30]. Briefly: for TEM, cells were seeded in a 6-well plate onto sapphire discs and grown for 72h with/without 0.5mM dbcAMP. After treatment, cells were fixed with 2.5% glutaraldehyde in 0.1M Na/K-phosphate, pH 7.4 for 1h

at 4°C, and kept in 0.1M Na/K-phosphate overnight at 4°C. Then, samples were post-fixed with 1% osmium tetroxide (OsO₄) in 0.1M Na/K-phosphate for 1h at 4°C. Afterwards, dehydration was performed in a graded series of ethanol starting at 70% followed by two changes in acetone and embedding in Epon. Sections 60-80 nm thick were stained with uranyl acetate and lead citrate and analysed in a transmission electron microscope (CM12, Philips, Eindhoven, The Netherlands) equipped with a CCD camera (Ultrascan 1000, Gatann Pleasanton, CA, USA) at an acceleration voltage of 100 kV.

For SEM, cells were grown on cover slips 10 mm in diameter (Mattek, Ashland, MA, USA). Following stimulation under the conditions described above, 2.5% glutaraldehyde in 0.1 M Na/K-phosphate, pH 7.4, was added to the warm culture medium. Then dishes were kept at 4°C for 1h, the medium was replaced by 0.1 M Na/K-phosphate, and stored at 4°C overnight. After post-fixation with 1% osmium tetroxide in 0.1M Na/K-phosphate, pH 7.4, at 4°C, cells were dehydrated in a graded series of ethanol starting at 10%, then critical-point dried (BAL-TEC, CPD 030, Balzers, Liechtenstein), coated with 4 nm platinum in a high vacuum sputtering device, (BAL-TEC SCD500), and examined in a SEM (Zeiss Supra 50 VP, Zeiss, Oberkochen, Germany) at an acceleration voltage of 5 kV using the secondary electron detector.

6.5 RNA isolation, reverse transcription (RT), semi-quantitative (TaqMan) PCR and data evaluation

Total RNA was isolated from primary stromal cells using TRIZOL[®]-Reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. The RNA content was measured with a NanoDrop 2000 UV-Vis Spectrophotometer[®] (Thermo Fisher Scientific, Reinach, CH). In order to remove potential genomic DNA contaminants resulting from the TRIZOL[®] protocol, RQ1 RNase-free DNase (Promega, Dübendorf, CH) was used following the

manufacturer's protocol. For complementary DNA (cDNA) synthesis, random hexamers were used as primers with RT reagents from Applied Biosystems, Foster City, CA, USA, following the manufacturer's protocol and as previously published [28, 31]. Reverse transcription was carried out in an Eppendorf Mastercycler[®] (Vaudaux-Eppendorf AG, Basel, CH) under the following conditions: 8 min at 21°C and 15 min at 42°C, after which the reaction was stopped by incubation for 5 min at 99°C. Semi-quantitative real time (TaqMan) PCR was performed in an automated fluorometer ABI PRISM[®] 7500 Sequence Detection System (Applied Biosystems) according to our previously described protocol [20, 32]. Fast Start Universal Probe Master (ROX[®]) (Roche Diagnostics AG, CH) was used with cDNA corresponding to 200ng of DNase-treated total RNA per sample. All reactions were run in duplicate. The following negative controls were applied: autoclaved water instead of cDNA and the so-called "RT minus" controls (samples which were not reverse transcribed), allowing exclusion of potential contaminants in reagents used for RT-PCR and confirming the accuracy of DNase-treatment, respectively. The semi-quantitation of target gene expression was carried out using three different endogenous reference genes (GAPDH, 18SrRNA and cyclophilin A). The list of primers and TaqMan Probes is presented in Table 1. The comparative CT method ($\Delta\Delta CT$ method) was applied according to the instructions of the ABI PRISM 7500 Sequence Detection System and as previously published [20, 32]: expression of each target gene relative to the reference genes, and normalised to the calibrator, *i.e.*, sample with the lowest amount of the respective target gene transcripts, was calculated. The average amounts of relative gene expression determined in relation to each of the reference genes for a sample were used as the normalization factor. The specificity of the selected PCR products was confirmed by sequencing (Microsynth, Balgach, CH). Canine-specific TaqMan Gene Expression Assays purchased from Applied Biosystems were used for: cyclophilin A (Prod. No. Cf03986523-gH), IGF-1 (Prod. No. Cf02627846_m1) and IGF-2 (Prod. No. Cf02647136_m1). Selected amplicons of each of the genes were sent for commercial sequencing (Microsynth). Statistical

analysis was performed using the GraphPad 3.06 software. One-way analysis of variance (ANOVA) for each of the treatment groups, *i.e.* 24h, 48h and 72h, was applied followed by the Tukey-Kramer Multiple Comparison Test. Numerical data are presented as the mean \pm standard deviation (SD).

6.6 Western Blot analysis

NET-2 lysis buffer (50mM Tris-HCl, PH 7.4, 300mM NaCl, 0.05% NP-40) containing 10 μ l/ml protease inhibitor cocktail (Sigma-Aldrich Chemie) was used to prepare whole cell lysates for the immunoblotting analyses which were performed as previously described [33]. Briefly: protein homogenates (20-30 μ g of total protein) were separated on 10% polyacrylamide gel (Bio-Rad Laboratories GmbH, Munich, Germany) and transferred onto methanol-activated polyvinylidene difluoride (PVDF) membranes (Bio-Rad) and probed with specific antibodies. Due to the limited availability of canine-specific and/or cross reacting antibodies suitable for Western blot analysis, detection of decidualisation markers using this method was restricted to the following target proteins that are significantly upregulated during decidualisation, and for which Western blot-suitable antisera were available: COX2 (PTGS2; using monoclonal mouse anti-rat PTGS2 IgG, clone 33, BD Pharmingen, Heidelberg, DE; dilution 1:500), and PTGES and PGFS/AKR1C3 (both guinea pig polyclonal, affinity-purified anti-dog, Eurogentec S.A., Seraing, Belgium; dilutions 1:25 and 1:1000, respectively; [33, 34]). The PVDF membranes were reblotted with monoclonal mouse anti-human GAPDH IgG (sc-47724) from Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA, at a dilution of 1:1000. Secondary antibodies were: goat anti-mouse IgG (W402B) from Promega (dilution 1:15,000) and anti-guinea pig IgG conjugated HRP from Sigma-Aldrich (A5545; 1:15,000). Detection of signals was performed using SuperSignal West chemiluminescent substrate (Life Technologies Europe B.V., Zug, CH) according to the manufacturer's protocol; visualization

of signals was done using the ChemiDoc XRS+ System and Image Lab Software, both from Bio-Rad. The optical density of bands was assessed using ImageJ software. In order to test for the effects of time and treatment on COX2 (PTGS2) and PTGES expression, an unpaired two-tailed Student's *t*-test was performed with the GraphPad 3.06 software program. Numerical data are presented as the mean \pm standard deviation (SD). Representative Western blots are shown.

7 Results

7.1 Characterisation and *in vitro* decidualisation of isolated canine uterine stromal cells

Primary stromal cells were isolated from uteri of dogs collected during the early dioestrus phase using enzymatic dissociation and using the differential adhesion time. Nearly 100% of cells stained positively for vimentin, thus verifying their mesenchymal origin (Figure 1). Only sporadically were epithelial cell-derived cytokeratin signals detected. The morphological changes observed during *in vitro* decidualisation were consistent with the acquired characteristics of decidual cells. The shape of cells changed from elongated and spindle-shaped to rounded (Figure 2A) when cells were incubated for 72h with 0.5 mM dbcAMP. Morphological changes observed during *in vitro* decidualisation of canine stromal cells were captured using time-lapse microscopy and are presented in the supplemental material (Supplement 1). Simultaneously with the morphological changes, the *in vitro* decidualised cells revealed an accelerated proliferation rate as determined by quantification of cell densities in time-lapse derived pictures at the selected time points of 0, 12, 24, 48 and 72h (Figure 2A). Whereas both dbcAMP-stimulated and control cell densities increased significantly over time ($P < 0.0001$), the density of decidualised cells significantly exceeded that of control cells by 48h and 72h ($P < 0.0002$ and $p < 0.0007$, respectively) (Figure 2B). Nearly 100% confluence was observed in decidualised cells by 72h.

Decidualised canine stromal cells displayed strong immunocytochemical signals for α SMA (Figure 3A). Its expression was compared with the expression and localization of α SMA in the pre-implantation canine uterus and in mid-term placenta. Prior to implantation, strong signals were localized in smooth muscle cells of the myometrium and tunica media and basement membrane of blood vessels, while weaker staining was observed in endometrial

stromal cells (Figure 3B). In addition to the vascular staining within the placenta, strong signals were localized in maternal decidual cells. (Figure 3C,D).

7.2 Subcellular morphological characterisation of canine decidual cells during *in vitro* decidualisation.

Transmission electron microscopic analysis (TEM) of control stromal cells revealed the presence of long, often branched mitochondria, frequently visible microfilaments and abundant cisternae of rough endoplasmic reticulum (Figure 4A-D). Cells undergoing *in vitro* decidualisation were characterized by higher abundance of Golgi apparatus which is responsible for protein sorting, modification and packaging for secretion, and consequently, greater amounts of secretory vesicles (Figure 4E-I). These cells also displayed higher numbers of small, round-shaped mitochondria. As revealed by scanning electron microscopic analysis (SEM), the increased secretory activity of decidualised cells was also clearly visible on cell surfaces, displaying higher amounts of secreted deposits than in non-treated controls (Figure 5A-F).

7.3 Expression of selected genes in canine uterine primary stromal cells during *in vitro* decidualisation

Table 1 presents a list of several selected genes, the expression of which was investigated at the mRNA level during dbcAMP-induced *in vitro* decidualisation of canine uterine stromal cells. Non-stimulated cells served as controls over the 72h time course of the study. Whereas the expression of IGF1R did not change significantly during the period of *in vitro* decidualisation, IGF1 was already strongly upregulated ($P < 0.05$) after 24h in response to 0.5 mM dbcAMP, while the expression of IGF2 was significantly elevated at 72h ($p < 0.05$) (Figure 6). The expression of PRLR increased significantly over controls within 24h,

responding to low and high dbcAMP concentrations ($P<0.05$, $P<0.01$ and $P<0.001$ for 0.1mM, 0.3mM and 0.5mM dbcAMP, respectively). This effect was also apparent in cells incubated for 48h ($P<0.001$ and $P<0.01$ for 0.3mM and 0.5mM dbcAMP, respectively). The highest PRLR response was noted at 72h in response to 0.3mM and 0.5mM dbcAMP, $P<0.001$ and $P<0.01$, respectively (Figure 6). In contrast, the expression of PRL mRNA remained low and frequently below the detection limit both in control and stimulated cells throughout the experimental time-course (not shown).

The evaluation of steroid hormone receptor expression revealed significantly ($P<0.05$) altered expression of ER α (ESR1) during *in vitro* decidualisation, presenting elevated mRNA levels after 48h and 72h in cells stimulated with 0.3mM or 0.5mM dbcAMP. The expression of PGR did not change significantly in decidualised cells ($P>0.05$) (Figure 6). Similarly, as for PRL, the expression of ER β (ESR2) mRNA remained mainly below the detection limit in all samples throughout the experiment (not shown).

Among the major members of the prostaglandin system, the expression of COX2 (PTGS2) displayed an expression pattern similar to that observed for IGF2 (Figure 7). It was, however, already significantly upregulated 24h after stimulation in response to 0.5mM dbcAMP ($P<0.01$). As for IGF2, its highest expression above controls was noted at 72h ($P<0.001$) in response to 0.5mM dbcAMP. Similarly, the expression of PGFS/AKR1C3 resembled that of IGF2 and was significantly increased at 72h in response to both 0.3mM and 0.5mM dbcAMP compared to the respective controls ($P<0.05$). The PGF2 α receptor PTGFR (FP) and PTGES presented similar expression levels (Figure 7). Thus, while displaying greater variations, their expression levels were increased ($P<0.05$) at 48h after stimulation. Especially for PTGES, this effect was obvious at 72h in response to the highest dbcAMP dosage ($P<0.05$). Contrasting with this was the expression of the respective PGE2 receptors, which either did not change significantly ($P>0.05$) for EP4 (PTGER4) or decreased significantly for EP2 (PTGER2), both in response to dbcAMP within the first 24h of incubation ($P>0.001$) when compared with the

respective control, as well as in all controls over the entire time course ($P < 0.001$ for 48h and $P < 0.01$ for 72h, respectively).

Similarly, as for EP4, the expression of PGT (SLCO2A1) remained unaffected ($P > 0.05$) during decidualisation. At the protein level, PTGS2 and PTGES expression profiles matched their respective mRNA levels, as shown in Figure 8 for 24h and 72h of stimulation. Thus, whereas COX2 (PTGS2) expression already increased significantly ($P < 0.01$) at 24h, both PTGS2 and PTGES were strongly upregulated at 72h ($P < 0.004$ and $p < 0.007$, respectively), compared with their respective controls. Although the canine-specific anti-PGFS/AKR1C3 antibody previously tested positively on canine uteri and placenta [33], within the current study the expression of PGFS/AKR1C3 protein remained below the detection limit (not shown).

8 Discussion

Decidualisation is a complex process, during which species displaying invasive types of placentation develop highly specialized maternal stroma-derived decidual cells. Their roles are in facilitating implantation and supporting embryo growth, *e.g.*, by secreting hormones, such as prolactin or placental lactogen, or cytokines involved in immunological protection of the foetus [35-37]. None of these functions, however, are known for the dog.

Although devoid of an anti-luteolytic mechanism that serves in other domestic animal species to prevent luteolysis, the canine uterus was recently shown to respond to the presence of pre-implantation, free-floating embryos by activating the expression of markers of ongoing, early decidualisation [15]. Following implantation and placentation, canine decidual cells eventually develop. These are the most important sensors of luteal P4 at the foeto-maternal interface in the dog [20]. Until now, the process of decidual cell formation in the dog was unknown, and neither was there an available cellular model allowing for detailed *in vitro* studies. Therefore, the present study investigated the basic capability of canine uterine stromal cells isolated from bitches during early dioestrus to decidualise *in vitro*. The dbcAMP-based protocol was chosen to compensate for our limited access to the tissue material which did not allow more detailed studies utilizing, *e.g.*, ovarian steroids. Nevertheless, dbcAMP has proved to be a potent stimulus mediating P4-driven decidualisation [23, 38]. Among other cAMP inducers capable of initializing decidualisation of uterine stromal cells are relaxin and gonadotropins [39, 40]. In line with this, responding to dbcAMP, canine stromal cells acquired the characteristics of decidual cells, reaching nearly 100% confluency at 72h. They changed morphologically, became rounded, exhibited accelerated proliferation rates and increased their metabolic activity, as concluded from the increased incidence of small mitochondria, elevated numbers of secretory follicles and increased deposits of extracellular matrix observed under scanning electron microscope. The composition and function of the

matrix components secreted by these cells in dogs needs to be determined. As a further sign of successful *in vitro* decidualisation, strong expression of α SMA was observed in these decidualised cells. Its expression is a typical sign of decidualisation known from other species including humans and is also an important marker used for detection of endometriotic lesions in women [41]. During the pre-implantation stage of pregnancy, the expression of α SMA in canine uterine stromal cells was only weak, indicating an early stage of uterine differentiation. Following placentation, it was localized and abundantly expressed in decidual cells of the *placenta materna*, which conformed to our *in vitro* findings.

At the biochemical and functional levels, the process of canine decidualisation still remains to be elucidated. In the present study, in order to validate the newly established *in vitro* model, the presence of several factors was investigated, the expression of which was recently described in early pregnant canine uterus and/or placenta [15, 20, 33, 34]. Some of them, such as IGF1 and IGF2 are among the most prominent decidualisation markers in other species [42, 43]. Others, like PGs, even if *per se* are not decidual cells markers, belong to strong inducers of the decidualisation process [23].

Thus, acting predominantly at the level of their common receptor IGFR1, both IGFs are mitogenic factors [42, 43]. They drive embryonic development and are important determinants of fetal and placental growth (reviewed in [44]). Their expression during *in vitro* decidualisation of canine stromal cells, as well as the expression of other factors investigated in the present study, was found to closely resemble their expression patterns evoked in early pregnant uteri in response to free-floating embryos [15]. Hence, both IGF1 and IGF2 responded positively to the applied decidualisation protocol. Whereas IGF1 turned out to be a quick responder displaying a significant dbcAMP dose-dependent response within 24h, IGF2 responded in a time-dependent manner with its highest expression observed only after 72h. Similar effects on IGF2 expression, becoming significant after 3 days of *in vitro* decidualisation, were observed in human endometrial stromal cells [38]. In contrast to human

and rodent models, however, in which PRL is one of the strongest decidualisation markers [38, 45, 46] involved in regulating endometrial secretory activity [47], canine decidualised cells exhibited very low expression of PRL. This, together with the concomitantly highly upregulated expression of PRLR observed in these cells, confirms our previous findings on the low expression of PRL in canine uteri which is frequently below detection limits [15], and points towards species-specific regulatory mechanisms. As part of these, it appears plausible that the strongly elevated levels of PRLR could compensate for low PRL expression, locally increasing its relative biological availability. This could also apply to the expression of ER α , which in contrast to ER β was strongly elevated during decidualisation, possibly compensating for the low availability of circulating oestradiol of luteal origin during canine gestation. This, indeed, strongly resembles the situation observed *in vivo* where higher ER α mRNA and protein levels were observed in early pregnant canine uteri compared to non-pregnant controls [15]. In this regard, interestingly, in naturally oestrogenized dogs, no additional supplementation with oestrogens is needed for establishment and maintenance of pregnancy. This was clearly shown by Concannon [48, 49] in experiments with dogs ovariectomized early in gestation (day 14 after the preovulatory LH surge), in which pregnancy was maintained only due to exogenous supplementation with P4. Although unaltered during *in vitro* and early *in vivo* decidualisation, the expression of PGR in canine decidual cells is a *sine qua non* condition for successful establishment and maintenance of canine pregnancy by preventing the premature luteolytic cascade [15, 20].

In human decidua, IGF1 was shown to be involved in regulation of arachidonic acid secretion, which is a common precursor for PG synthesis [50]. Similarly, in our experiments the *in vitro* expression of PG synthases closely paralleled the expression of IGFs. The functional interrelationship between these entities remains to be elucidated for the dog.

The versatile functions of PGs include the cAMP/PKA pathway-mediated stimulatory effects of PGE2 in further enhancing P4 and cAMP-induced decidualisation in humans [23]. Also,

immunosuppressive but embryo-protective effects of human decidua-derived PGE2 were proposed [51]. Furthermore, in addition to the species-specific function of oestrogens, both, embryo- and luteo-protective roles of endometrial PGE2 have been proposed for pigs (see review [52]). Since all of the above functions could also apply to the dog, the modulatory effect of decidualisation on the expression of PG family members in canine stromal cells is noteworthy. PGs, and especially PGE2, are among the most important luteotrophic factors in the dog [27, 53-55]. Together with IGF2, abundant expression of PTGES was observed in hatched canine pre-implantation embryos and the pre-implantation uterus [15]. Interestingly, during placentation PGs are more strongly represented in the foetal compartments [20, 33, 34]. However, the increased expression shown herein of COX2 and PTGES transcripts, but not those of PGFS/AKR1C3, was clearly mirrored at the protein level. Thus, in addition to the foetal trophoblast, decidual cells appear to be an important potential source of canine placental PGs. By expressing the respective PG receptors and PGT (SLCO2A1), they are most likely involved in paracrine and autocrine local regulatory mechanisms.

Taking into account the indispensable function of decidual cells in ensuring proper uterine secretory activity, as well as their roles during the establishment and maintenance of canine pregnancy, herein a valuable *in vitro* model was established showing the basic capability of canine uterine stromal cells to decidualise *in vitro*. The morphological and biochemical characteristics of decidual cells were acquired during this process, providing evidence that canine decidualisation can be effectively modeled *in vitro*. Nevertheless, further investigations are needed to dissect the underlying molecular and endocrine mechanisms involved. In particular, the role of upstream operating regulatory mechanisms responsible for initializing the cAMP-dependent decidualisation cascade, needs to be addressed.

9 Figures and Legends


9.1 Table 1

List of primers and TaqMan Probes used for the semi-quantitative RT-PCR.

Primer	Accession numbers	Primer sequence	Product length (bp)
<i>GAPDH</i>	AB028142	Forward: 5'-GCT GCC AAA TAT GAC GAC ATC A-3' Reverse: 5'-GTA GCC CAG GAT GCC TTT GAG-3' TaqMan Probe: 5'-TCC CTC CGA TGC CTG CTT CAC TAC CTT-3'	75
<i>18SrRNA</i>	FJ797658	Forward: 5'-GTC GCT CGC TCC TCT CCT ACT-3' Reverse: 5'-GGC TGA CCG GGT TGG TTT-3' TaqMan Probe: 5'-ACA TGC CGA CGG GCG CTG AC-3'	125
<i>IGFR1</i>	XM545828	Forward: 5'-GGA CGT TGA GCC TGG CAT T-3' Reverse: 5'-CAC TCT TAG CCC CAC GGA TGT-3' TaqMan Probe: 5'-AGC CCT GGA CGC AGT ATG CGG-3'	119
<i>PRLR</i>	HQ267784	Forward: 5'-GGA TCT TTG TGG CCG TTC TTT-3' Reverse: 5'-AAG GAT GCA GGT CAC CAT GCT AT-3' TaqMan Probe: 5'-ATT ATG GTC GTA GCA GTG GCT TTG AAA GGC-3'	92
<i>PRL</i>	NM_001008275	Forward: 5'-CAAGCC CAA CAG ATC CAC CAT-3' Reverse: 5'-ATC CCC CGC ACT TCT GTG A-3' TaqMan Probe: 5'-CTG AGG GTG CTG CGC TCC TGG-3'	104
<i>PGR</i>	NM_001003074	Forward: 5'-CGA GTC ATT ACC TCA GAA GAT TTG TTT-3' Reverse: 5'-CTT CCA TTG CCC TTT TAA AGA AGA-3' TaqMan Probe: 5'-AAG CAT CAG GCT GTC ATT ATG GTG TCC TAA CTT-3'	113
<i>ESR1</i> (<i>ERα</i>)	XM533454	Forward: 5'-CCC ATG GAG GAG ACA AAC CA-3' Reverse: 5'-CCC TGC CTC CTC GGT GAT ATA-3' TaqMan Probe: 5'-CAC GGG CCC AAC TTC ATC ACA TTC C-3'	93
<i>ESR2</i> (<i>ERβ</i>)	XM861041	Forward: 5'-CCC AGC ACG CCC TTC A-3' Reverse: 5'-AAT CAT ATG CAC GAG TTC CTT GTC-3' TaqMan Probe: 5'-CCT CCA TGA TGA TGT CCC TGA CC-3'	78
<i>PTGS2</i>	HQ110882	Forward: 5'-GGA GCA TAA CAG AGT GTG TGA TGT G-3' Reverse: 5'-AAG TAT TAG CCT GCT CGT CTG GAA T-3' TaqMan Probe: 5'-CGC TCA TCA TCC CAT TCT GGG TGC-3'	87
<i>AKR1C3</i>	NM_001012344	Forward: 5'-AGG GCT TGC CAA GTC TAT TGG-3' Reverse: 5'-GCC TTG GCT TGC TCA GGA T-3' TaqMan Probe: 5'-TCC AAC TTT AAC CGC AGG CAG CTG G-3'	74
<i>PTGES</i>	NM_001122854	Forward: 5'-GTC CTG GCG CTG GTG AGT-3' Reverse: 5'-ATG ACA GCC ACC ACG TAC ATC T-3' TaqMan Probe: 5'-TCC CAG CCT TCC TGC TCT GCA GC-3'	89
<i>PTGFR (FP)</i>	NM_001048097	Forward: 5'-ACC AGT CGA ACA TCC TTT GCA-3' Reverse: 5'-GGC CAT CAC ACT GCC TAG AAA-3' TaqMan Probe: 5'-CAT GGT GTT CTC CGG TCT GTG CCC-3'	86
<i>PTGER2</i>	AF075602	Forward: 5'-CAC CCT GCT GCT GCT TCT C-3' Reverse: 5'-CGG TGC ATG CGG ATG AG-3' TaqMan Probe: 5'-TGC TCG CCT GCA ACT TTC AGC GTC-3'	78
<i>PTGER4</i>	NM_001003054	Forward: 5'-AAA TCA GCA AAA ACC CAG ACT TG-3' Reverse: 5'-GCA CGG TCT TCC GCA GAA-3' TaqMan Probe: 5'-ATC CGA ATT GCT GCT GTG AAC CCT ATC C-3'	96
<i>SLCO2A1</i> (<i>PGT</i>)	NM_001011558	Forward: 5'-TGC AGC ACT AGG AAT GCT GTT C-3' Reverse: 5'-GGG CGC AGA GAA TCA TGG A-3' TaqMan Probe: 5'-TCT GCA AAC CAT TCC CCG CGT G-3'	116


9.2 Figure 1

Canine primary uterine stromal cells. (A) A high confluence monolayer is presented. (B) Vimentin (green) and cytokeratin (red, inset in B) immunofluorescence was performed in order to confirm the mesenchymal origin of uterine stromal cells. There is no or only weak staining for cytokeratin visible indicating high purity of isolated stromal cells. C and D present merged pictures at lower and higher magnifications with cells stained for vimentin and cytokeratin; nuclear staining was achieved with 4',6-Diamidino-2-phenylindole (DAPI).




9.3 Figure 2

Morphological appearance of canine uterine stromal cells during *in vitro* decidualisation (A; presented at 100x magnification). Quantitation of cell density: percentage (%) of surface determined under live cell imaging conditions (B; average from three independent experiments). One-way ANOVA followed by Dunnett's Multiple Comparison Test was applied to test the effects of time on cell density in all control or treated samples ((*) indicates $P < 0.0001$). Student's *t*-test was applied to test the effect of treatment on cell density: (**) indicates $P < 0.0002$, (***) indicates $P < 0.0007$.




9.4 Figure 3

Immunostaining against smooth muscle alpha actin (α SMA). (A) Decidualised canine stromal cells expressing α SMA (green; nuclear staining was achieved by DAPI). (B) Immunohistochemical localization of α SMA in canine pre-implantation uterus (solid arrows = blood vessels; open arrows = smooth muscle cells of myometrium, open arrowheads = uterine stromal cells, solid arrowheads = uterine glands). (C, D) Immunohistochemical localization of α SMA in canine mid-term placenta (solid arrows = large maternal vessels; open arrows = decidual cells).




9.5 Figure 4

Transmission electron microscopy (TEM) analysis of canine uterine primary stromal cells during *in vitro* decidualisation. (A-D) control cells at 72h of culture (solid arrowheads = nucleus, open arrows = mitochondria, solid arrows = microfilaments, open arrowheads = cisternae of rough endoplasmic reticulum). (E-I) *in vitro* decidualised cells, after 72h treatment with 0.5 mM dbcAMP (solid arrowheads = large secretory vesicles, open arrows = mitochondria, solid arrows = Golgi apparatus, open arrowheads = cisternae of rough endoplasmic reticulum).




9.6 Figure 5

Scanning electron microscopy (SEM) imaging of canine uterine primary stromal cells during *in vitro* decidualisation. (A-C) control cells at 72h of culture, (D-F) *in vitro* decidualised cells, after 72h treatment with 0.5 mM dbcAMP. Increased secretory activity of decidualised cells is visible on cell surface, as characterized by larger amounts of secreted deposits (open arrows in D-F).




9.7 Figure 6

Expression of insulin-like growth factor (IGF1), IGF2, IGF-receptor 1 (IGF1R), prolactin receptor (PRLr), progesterone receptor (PGR) and oestrogen receptor alpha ($ER\alpha$, ESR1), as determined by Real Time (TaqMan) PCR. Canine primary stromal cells were cultured for 24, 48 and 72h in the presence of increasing dbcAMP concentrations. One-way ANOVA (24 h in IGF1 $P < 0.01$, 72h in IGF1 $P < 0.02$; 72h in IGF2 $P < 0.03$; 24 h in PRLR $P < 0.0003$, 48h in PRLR $P < 0.0001$, 72h in PRLR $P < 0.001$; 48 h in $ER\alpha$ $P < 0.02$, 72h in $ER\alpha$ $P < 0.004$), followed by the Tukey-Kramer Multiple Comparison Test was applied; all samples were compared against the non-treated control in each group. Different letters indicate $P < 0.05$. Numerical data are presented as the mean \pm standard deviation (SD).




9.8 Figure 7

Expression of cyclooxygenase 2 (COX2, PTGS2), PGE2-synthase (PGES, PTGES), PGE2 receptors: EP4 (PTGER4) and EP2 (PTGER2), PGF2 α -synthase (PGFS/AKR1C3), PGF2 α receptor (PTGFR, FP) and PG-transporter (PGT, SLCO2A1), as determined by Real Time (TaqMan) PCR. Canine primary stromal cells were cultured for 24, 48 and 72h in the presence of increasing dbcAMP concentrations. One-way ANOVA was applied (24 h in COX2 $P < 0.005$, 48h in COX2 $P < 0.01$, 72h in COX2 $P < 0.0001$; 48 h in PGES $P < 0.01$, 72h in PGES $P < 0.03$; 24h in EP2 $P < 0.0001$, 72h in PGFS $P < 0.03$, 48h in FP $P < 0.03$, 72h in FP $P < 0.01$), followed by the Tukey-Kramer Multiple Comparison Test; all samples were compared against the non-treated control in each group. Different letters indicate $P < 0.05$. Numerical data are presented as the mean \pm standard deviation (SD).



9.9 Figure 8

Expression of cyclooxygenase 2 (COX2, PTGS2) and PGE2-synthase (PGES, PTGES) as determined by Western blot analysis. Following stimulation, cells were collected and homogenized and 20 μ g of the lysate was used for Western blots. The averaged standardized optical density (SOD) is shown. GAPDH was used for loading control. The cell culture experiments were performed independently at least three times using cells isolated from different animals. Representative Western blots are shown. Student's *t*-test was applied to test the effect of treatment on protein expression: in (A) (*) indicates $P < 0.01$, (**) indicates $P < 0.004$, (***) indicates $P < 0.0001$; in B (*) indicates $P < 0.008$.



9.10 Supplement 1

Time-lapse videos of *in vitro* decidualising canine uterine stromal cells compared with untreated controls. Both representative videos were taken simultaneously during one experiment over 72h (movies are displayed at 100x magnification).

(for supplemental file see CD ROM)

10 Funding

Swiss National Science Foundation (SNSF) research grant number 31003A_140947.

11 References

1. Gellersen B, Brosens IA, Brosens JJ. Decidualization of the human endometrium: mechanisms, functions, and clinical perspectives. *Semin Reprod Med.* 2007; 25 (6):445-53.
2. Aplin J. Maternal influences on placental development. *Semin Cell Dev Biol.* 2000; 11 (2):115-25.
3. Ramathal CY, Bagchi IC, Taylor RN, Bagchi MK. Endometrial decidualization: of mice and men. *Semin Reprod Med.* 2010; 28 (1):17-26.
4. Frolova A, Flessner L, Chi M, Kim ST, Foyouzi-Yousefi N, Moley KH. Facilitative glucose transporter type 1 is differentially regulated by progesterone and estrogen in murine and human endometrial stromal cells. *Endocrinology.* 2009; 150(3):1512-20.
5. Sharkey AM, Smith SK. The endometrium as a cause of implantation failure. *Best Pract Res Clin Obstet Gynaecol.* 2003; 17 (2):289-307.
6. Jauniaux E, Jurkovic D. Placenta accreta: pathogenesis of a 20th century iatrogenic uterine disease. *Placenta.* 2012; 33 (4):244-251.
7. Al-Bassam MA, Thomson RG, O'Donnell L. Involution abnormalities in the postpartum uterus of the bitch. *Veter pathol.* 1981; 18 (2):208-18.
8. Nishiyama T, Tsumagari S, Ito M, Kimura J, Watanabe G, Taya K, et al.. Immunohistochemical study of steroidogenic enzymes in the ovary and placenta during pregnancy in the dog. *Anat Histol Embryol.* 1999; 28 (2):125-9.
9. Hoffmann B, Hoveler R, Nohr B, Hasan SH. Investigations on hormonal changes around parturition in the dog and the occurrence of pregnancy-specific non conjugated oestrogens. *Exp Clin Endocrinol.* 1994; 102 (3):185-9.
10. Concannon PW, McCann JP, Temple M. Biology and endocrinology of ovulation, pregnancy and parturition in the dog. *J Reprod Fertil Suppl* 1989, 39:3-25.
11. Hoffmann B, Hoveler R, Hasan SH, Failing K. Ovarian and pituitary function in dogs after hysterectomy. *J Reprod Fertil.* 1992; 96 (2):837-45.
12. Thatcher WW, Bartol FF, Knickerbocker JJ, Curl JS, Wolfenson D, Bazer FW, et al.. Maternal recognition of pregnancy in cattle. *J Dairy Sci.* 1984; 67 (11):2797-811.
13. Bazer FW, Thatcher WW, Hansen PJ, Mirando MA, Ott TL, Plante C. Physiological mechanisms of pregnancy recognition in ruminants. *J Reprod Fertil Suppl.* 1991; 43:39-47.

14. Perry JS, Heap RB, Amoroso EC. Steroid hormone production by pig blastocysts. *Nature*. 1973; 245 (54):45-47.
15. Kautz E, Gram A, Aslan S, Ay SS, Selcuk M, Kanca H, et al.. Expression of genes involved in the embryo-maternal interaction in the early-pregnant canine uterus. *Reproduction*. 2014; 147 (5):703-17.
16. Frasier J, Gaspar CA, Donnelly KM, Gibori G, Fazleabas AT. Expression of prolactin and its receptor in the baboon uterus during the menstrual cycle and pregnancy. *J Clin Endocrinol Metab*. 1999; 84 (9):3344-50.
17. Goffin V, Binart N, Touraine P, Kelly PA. Prolactin: the new biology of an old hormone. *Annu Rev Physiol*. 2002; 64:47-67.
18. Amoroso EC. Placentation. In: Parkers AS (ed.) *Marshall's physiology of reproduction*. Longmans Green, London 1952; 127-316
19. Kehrer A. [Chorionic development and structure of the placenta zonaria in the cat, dog and fox (author's transl)]. *Z Anat Entwicklungsgesch*. 1973; 143:25-42.
20. Kowalewski MP, Beceriklisoy HB, Pfarrer C, Aslan S, Kindahl H, Kucukaslan I, et al.. Canine placenta: a source of prepartal prostaglandins during normal and antiprogesterin-induced parturition. *Reproduction*. 2010; 139 (3):655-64.
21. Vermeirsch H, Simoens P, Lauwers H. Immunohistochemical detection of the estrogen receptor-alpha and progesterone receptor in the canine pregnant uterus and placental labyrinth. *Anat Rec* 2000;. 260; (1):42-50.
22. Gram A, Boos A, Kowalewski MP. Uterine and placental expression of canine oxytocin receptor during pregnancy and normal and induced parturition. *Reprod Domest Anim*. 2014; 49 Suppl (2), 2:41-9.
23. Brar AK, Frank GR, Kessler CA, Cedars MI, Handwerger S. Progesterone-dependent decidualization of the human endometrium is mediated by cAMP. *Endocrine*. 1997; 6 (3):301-7.
24. Gellersen B, Brosens J. Cyclic AMP and progesterone receptor cross-talk in human endometrium: a decidualizing affair. *J Endocrinol*. 2003; 178 (3):357-72.
25. Kim JJ, Fazleabas AT. Uterine receptivity and implantation: the regulation and action of insulin-like growth factor binding protein-1 (IGFBP-1), HOXA10 and forkhead transcription factor-1 (FOXO-1) in the baboon endometrium. *Reprod Biol Endocrin*. 2004; 2:34.
26. Tamura I, Asada H, Maekawa R, Tanabe M, Lee L, Taketani T, et al.. Induction of IGFBP-1 expression by cAMP is associated with histone acetylation status of the

- promoter region in human endometrial stromal cells. *Endocrinology*. 2012; 153 (11):5612-21.
27. Kowalewski MP, Fox B, Gram A, Boos A, Reichler I. Prostaglandin E2 functions as a luteotrophic factor in the dog. *Reproduction*. 2013; 145 (3):213-26.
 28. Kowalewski MP, Schuler G, Taubert A, Engel E, Hoffmann B. Expression of cyclooxygenase 1 and 2 in the canine corpus luteum during diestrus. *Theriogenology*. 2006; 66 (6-7):1423-30.
 29. Kowalewski MP, Mason JI, Howie AF, Morley SD, Schuler G, Hoffmann B. Characterization of the canine 3beta-hydroxysteroid dehydrogenase and its expression in the corpus luteum during diestrus. *J Steroid Biochem Mol Biol*. 2006; 101 (4-5):254-62.
 30. Wild P. Electron microscopy of viruses and virus-cell interactions. *Method Cell Biol*. 2008; 88:497-524.
 31. Kowalewski MP, Michel E, Gram A, Boos A, Guscetti F, Hoffmann B, et al.. Luteal and placental function in the bitch: spatio-temporal changes in prolactin receptor (PRLr) expression at dioestrus, pregnancy and normal and induced parturition. *Reprod Biol Endocrin*. 2011; 9:109.
 32. Kowalewski MP, Meyer A, Hoffmann B, Aslan S, Boos A. Expression and functional implications of peroxisome proliferator-activated receptor gamma (PPARgamma) in canine reproductive tissues during normal pregnancy and parturition and at antiprogestin induced abortion. *Theriogenology*. 2011; 75 (5):877-86.
 33. Gram A, Buechler U, Boos A, Hoffmann B, Kowalewski MP. Biosynthesis and degradation of canine placental prostaglandins: prepartum changes in expression and function of prostaglandin F2alpha-synthase (PGFS, AKR1C3) and 15-hydroxyprostaglandin dehydrogenase (HPGD). *Biol Reprod*. 2013a; 89 (1):2.
 34. Gram A, Fox B, Buchler U, Boos A, Hoffmann B, Kowalewski MP. Canine placental prostaglandin E2 synthase: expression, localization, and biological functions in providing substrates for prepartum PGF2alpha synthesis. *Biol Reprod*. 2014; 91 (6):154.
 35. Alam SM, Konno T, Dai G, Lu L, Wang D, Dunmore JH, et al.. A uterine decidual cell cytokine ensures pregnancy-dependent adaptations to a physiological stressor. *Development*. 2007; 134 (2):407-15.

36. Dudley DJ, Trautman MS, Araneo BA, Edwin SS, Mitchell MD. Decidual cell biosynthesis of interleukin-6: regulation by inflammatory cytokines. *J Clin Endocrinol Metab.* 1992; 74 (4):884-89.
37. Golander A, Hurley T, Barrett J, Hizi A, Handwerger S. Prolactin synthesis by human chorion-decidual tissue: a possible source of prolactin in the amniotic fluid. *Science.* 1978; 202 (4365):311-13.
38. Ganeff C, Chatel G, Munaut C, Franken F, Foidart JM, Winkler R. The IGF system in in-vitro human decidualization. *Mol Hum Reprod.* 2009, 15 (1):27-38.
39. Lane B, Oxberry W, Mazella J, Tseng L. Decidualization of human endometrial stromal cells in vitro: effects of progestin and relaxin on the ultrastructure and production of decidual secretory proteins. *Hum Reprod.* 1994; 9 (2):259-66.
40. Tang B, Gurpide E. Direct effect of gonadotropins on decidualization of human endometrial stroma cells. *J Steroid Biochem Mol Biol,* 1993; 47 (1-6):115-21.
41. Anaf V, Simon P, Fayt I, Noel J. Smooth muscles are frequent components of endometriotic lesions. *Hum Reprod.* 2000; 15 (4):767-71.
42. Wang HS, Chard T. IGFs and IGF-binding proteins in the regulation of human ovarian and endometrial function. *J Endocrinol.* 1999; 161 (1):1-13.
43. Yu J, Wu J, Bagchi IC, Bagchi MK, Sidell N, Taylor RN. Disruption of gap junctions reduces biomarkers of decidualization and angiogenesis and increases inflammatory mediators in human endometrial stromal cell cultures. *Mol Cell Endocrinol.* 2011; 344 (1-2):25-34.
44. Wathes DC, Reynolds TS, Robinson RS, Stevenson KR. Role of the insulin-like growth factor system in uterine function and placental development in ruminants. *J Dairy Sci.* 1998; 81 (6):1778-89.
45. Tseng L, Mazella J. Prolactin and its receptor in human endometrium. *Semin Reprod Med.* 1999; 17 (1):23-27.
46. Prigent-Tessier A, Tessier C, Hirosawa-Takamori M, Boyer C, Ferguson-Gottschall S, Gibori G. Rat decidual prolactin. Identification, molecular cloning, and characterization. *J Biol Chem* 1999, 274 (53):37982-89.
47. Jabbour HN, Critchley HO, Boddy SC. Expression of functional prolactin receptors in nonpregnant human endometrium: janus kinase-2, signal transducer and activator of transcription-1 (STAT1), and STAT5 proteins are phosphorylated after stimulation with prolactin. *J Clin Endocrinol Meta.* 1998; 83 (7):2545-53.

48. Concannon P, Tsutsui T, Shille V. Embryo development, hormonal requirements and maternal responses during canine pregnancy. *J Reprod Fertil Suppl* 2001, 57:169-79.
49. Concannon PW. Endocrinologic control of normal canine ovarian function. *Reprod Domest Anim*. 2009; 44 Suppl 2:3-15.
50. Handwerger S, Markoff E, Richards R. Regulation of the synthesis and release of decidual prolactin by placental and autocrine/paracrine factors. *Placenta*. 1991; 12 (2):121-30.
51. Parhar RS, Yagel S, Lala PK. PGE₂-mediated immunosuppression by first trimester human decidual cells blocks activation of maternal leukocytes in the decidua with potential anti-trophoblast activity. *Cell Immunol*. 1989; 120 (1):61-74.
52. Ziecik AJ, Waclawik A, Kaczmarek MM, Blitek A, Jalali BM, Andronowska A. Mechanisms for the establishment of pregnancy in the pig. *Reprod Domest Anim*. 2011; 46 Suppl 3:31-41.
53. Kowalewski MP, Mutembei HM, Hoffmann B. Canine prostaglandin E₂ synthase (PGES) and its receptors (EP₂ and EP₄): expression in the corpus luteum during dioestrus. *Anim Reprod Science*. 2008; 109 (1-4):319-29.
54. Kowalewski MP, Beceriklisoy HB, Aslan S, Agaoglu AR, Hoffmann B. Time related changes in luteal prostaglandin synthesis and steroidogenic capacity during pregnancy, normal and antiprogesterin induced luteolysis in the bitch. *Anim Reprod Sci*. 2009; 116 (1-2):129-38.
55. Kowalewski MP, Ihle S, Siemieniuch MJ, Gram A, Boos A, Zdunczyk S, et al.. Formation of the early canine CL and the role of prostaglandin E₂ (PGE₂) in regulation of its function: An in vivo approach. *Theriogenology*. 2015; 83 (6):1038-47.

12 Acknowledgements

I would like to express my sincere appreciation to everybody who, in one way or another, contributed to my thesis.

My greatest appreciation and sincere thanks go to my supervisor, Prof. Mariusz P. Kowalewski, PhD for his supervision, patient guidance, endless encouragement, suggestions and useful critiques of this research work.

I would also like to express my sincere gratitude to the director of the Institute of Veterinary Anatomy, Prof. Dr. Alois Boos for his support and continuous encouragement.

Prof Dr. Jan Twardoń and Dr. Renata Nowaczyk have always been very supportive during my study for which I am most thankful.

I would also like to express my appreciation to the technicians of the laboratory of the Institute of Veterinary Anatomy, Elisabeth Hoegger, Elisabeth Schraner and Stefanie Ihle for their help and support.

I would also like to extend my sincere thanks to colleagues from Institute of Veterinary Anatomy for their help, kindness and friendship especially to Stefanie Ihle, Sophie Latter, Jeannette Attiger and Aykut Gram.

Last, but not least, I would like to express my deep gratitude to my parents Dorota and Stanisław Kautz, and to my sister Anna Kautz as well as to my boyfriend Carsten Roth for their understanding, support, encouragement, endless love and endless sacrifices during my study.

13 Curriculum vitae

Vorname, Name: Ewa, Kautz
Geburtsdatum: 06.03.1987
Geburtsort: Gostyń, Polen
Nationalität: Polnisch

09/1994 – 06/2000 Primarschule: Szkoła Podstawowa nr 3 w Gostyniu, Gostyń, Polen
09/2000 – 06/2003 Gymnasium: Gimnazjum nr 2 w Gostyniu, Gostyń, Polen
09/2003 – 06/2006 Zespół Szkół Ogólnokształcących w Gostyniu, Gostyń, Polen;
Abschluss mit Matura
10/2006 – 05/2012 Studium der Veterinärmedizin, Uniwersytet Przyrodniczy we
Wrocławiu, Wrocław, Polen
09/2012 – 03/2015 Anfertigung der Dissertation
unter Leitung von Prof. Dr. Mariusz P. Kowalewski
am Veterinär-Anatomischen Institut
der Vetsuisse-Fakultät Universität Zürich
Direktor Prof. Dr. Alois Boss